arXiv:1607.02929v1 [math-ph] 11 Jul 2016

QUASI-EXACTLY SOLVABLE SYMMETRIZED QUARTIC AND SEXTIC POLYNOMIAL OSCILLATORS

C. QUESNE

Physique Nucléaire Théorique et Physique Mathématique,

Université Libre de Bruxelles, Campus de la Plaine CP229,

Boulevard du Triomphe, B-1050 Brussels, Belgium

cquesne@ulb.ac.be

Abstract

The symmetrized quartic polynomial oscillator is shown to admit an sl(2,$\mathbb{R}$) algebraization. Some simple quasi-exactly solvable (QES) solutions are exhibited. A new symmetrized sextic polynomial oscillator is introduced and proved to be QES by explicitly deriving some exact, closed-form solutions by resorting to the functional Bethe ansatz method. Such polynomial oscillators include two categories of QES potentials: the first one containing the well-known analytic sextic potentials as a subset, and the second one of novel potentials with no counterpart in such a class.

Running head: Symmetrized polynomial oscillators

Keywords: quantum mechanics; quasi-exact solvability; polynomial oscillators

PACS Nos.: 03.65.Fd, 03.65.Ge

1 Introduction

Quasi-exactly solvable (QES) Schrödinger equations occupy an intermediate place between exactly-solvable and non-solvable ones. Such equations, which have widespread applications in physics, chemistry, and engineering, are characterized by the fact that a (finite) part of their spectrum can be determined algebraically if the potential parameters satisfy certain constraints, while the remaining part remains unknown. Among the procedures used to construct them, one may quote the recursion relation method, which is rather hard to perform since one has to deal with three- or more-step recursion relations (see, e.g., [1]). Two easier approaches are the one based on an sl(2,$\mathbb{R}$) algebraization [2, 3, 4, 5] and that relying on the functional Bethe ansatz method [6, 7, 8].

The most prominent example of QES systems is the sextic polynomial oscillator [9, 10]. In contrast, the non-QES nature of the simpler and phenomenologically more important quartic polynomial oscillator [3] has always been surprising. In a recent work inspired by the well-known example of the square well, Znojil proposed a resolution of this paradox by allowing an analyticity-violating symmetrization of the quartic polynomial oscillator and he proved that the resulting potential is QES by using the recursion relation approach [11].

This has emphasized the interest of splitting the real line of coordinates into subintervals and considering potentials that admit different definitions inside these subintervals, while remaining continuous on the whole real line [12, 13, 14]. Such a procedure indeed opens the possibility of constructing new ES or QES potentials.

The purpose of the present work is twofold: first to re-examine the symmetrized quartic polynomial oscillator in the light of the sl(2,$\mathbb{R}$) Lie algebraic method, and second to propose a new symmetrized sextic one, while providing some explicit solutions of both of them.

2 QES Symmetrized Quartic Polynomial Oscillator

As in [11], let us consider the symmetrized quartic polynomial oscillator

$$V(x) = x^4 - s|x|^3 + rx^2 - q|x| = \begin{cases} x^4 + sx^3 + rx^2 + qx & \text{if } x < 0, \\ x^4 - sx^3 + rx^2 - qx & \text{if } x > 0, \end{cases} \tag{2.1}$$

which is non-analytic and defined separately on the two semi-infinite intervals $x < 0$ and $x > 0$. Here q, r, and s are three real constants. The corresponding Schrödinger equation

$$\left(-\frac{d^2}{dx^2} + V(x)\right)\psi(x) = E\psi(x) \tag{2.2}$$

may be considered separately on these two intervals. Physically acceptable wavefunctions should tend to zero at infinity and be continuous everywhere, as well as their first derivative. Since the potential is symmetric under parity, they must also be either symmetric or antisymmetric. We may therefore solve Eq. (2.2) in one of the intervals, e.g., $x < 0$, then impose that in the other interval $x > 0$, $\psi(x) = \epsilon\psi(-x)$ with $\epsilon = +1$ or $\epsilon = -1$ according to whether ψ is even or odd.

QES solutions of (2.2) may be obtained by assuming that $\psi(x)$ takes the form

$$\psi_n^{(\epsilon)}(x) = e^{W(x)}\phi_n^{(\epsilon)}(x), \qquad W(x) = -\tfrac{1}{3}|x|^3 + ax^2 - b|x|, \tag{2.3}$$

with

$$\phi_n^{(\epsilon)} = \begin{cases} \sum_{k=0}^{n} v_k x^k & \text{if } x < 0, \\ \epsilon \sum_{k=0}^{n} (-1)^k v_k x^k & \text{if } x > 0. \end{cases} \tag{2.4}$$

Here a, b are two real constants, and v_k ($k = 0, 1, \ldots, n$) are some expansion coefficients to be determined. Such functions have the right behaviour for $x \to \pm\infty$. On the other hand, the continuity of $\psi_n^{(\epsilon)}(x)$ and of its derivative at the origin imposes the additional condition

$$\begin{aligned} v_1 + bv_0 &= 0 \qquad \text{if } \epsilon = +1, \\ v_0 &= 0 \qquad \text{if } \epsilon = -1. \end{aligned} \tag{2.5}$$

Inserting Eq. (2.3) in (2.2) and setting

$$s = 4a \qquad \text{and} \qquad r = 4a^2 + 2b \tag{2.6}$$

in order to eliminate some high-order terms, we arrive at the differential equation

$$\left(-\frac{d^2}{dx^2}-2(x^2+2ax+b)\frac{d}{dx}+(q-4ab-2)x-b^2-2a\right)\phi_n(x)=E_n\phi_n(x) \quad (2.7)$$

on the interval $x<0$. This is a (triconfluent) Heun equation [15]. Hence, provided we also set

$$q=4ab+2n+2, \quad (2.8)$$

the second-order differential operator h on its left-hand side can be written as a second-degree polynomial [5]

$$h=-J_n^-J_n^--2J_n^+-4aJ_n^0-2bJ_n^--2(n+1)a-b^2 \quad (2.9)$$

in the sl(2,$\mathbb{R}$) generators

$$J_n^+=x^2\frac{d}{dx}-nx, \qquad J_n^0=x\frac{d}{dx}-\frac{n}{2}, \qquad J_n^-=\frac{d}{dx} \quad (2.10)$$

acting in the $(n+1)$-dimensional representation in the space of polynomials of order not higher than n, $\mathcal{P}_{n+1}=\langle 1,x,x^2,\ldots,x^n\rangle$. On using the relations $J_n^+x^k=(k-n)x^{k+1}$, $J_n^0x^k=(k-\frac{n}{2})x^k$, $J_n^-x^k=kx^{k-1}$, it is straightforward to explicitly write the matrix of h in the space $\mathcal{P}_{n+1}$. Its diagonalization, followed by the imposition of constraint (2.5), will then provide the admissible energies $E_n^{(\epsilon)}$ and wavefunctions $\psi_n^{(\epsilon)}(x)$.

In [11], Znojil solved the rather complicated $n=2$, $\epsilon=+1$ case by the recursion relation method. In the remainder of this section, we plan to exhibit some simpler examples, which can be directly derived by using (2.7)–(2.9).

For $n=0$, we obtain $E_0=-2a-b^2$ and $\phi_0(x)=1$ on $x<0$. Since $v_0=1$ and $v_1=0$, according to (2.5), the resulting wavefunction on the whole real line cannot be odd, but it can be even if the condition $b=0$ is satisfied. We conclude that the potential

$$V(x)=x^4-4a|x|^3+4a^2x^2-2|x| \quad (2.11)$$

has an energy eigenvalue $E_0^{(+)}$ and associated wavefunction $\psi_0^{(+)}(x)$, given by

$$E_0^{(+)}=-2a, \qquad \psi_0^{(+)}(x)=e^{-\frac{1}{3}|x|^3+ax^2}, \quad (2.12)$$

corresponding to a ground state.

Turning now ourselves to the $n = 1$ case, we arrive at a second-degree equation for E_1, whose solutions $E_{1\pm} = -4a - b^2 \pm 2\sqrt{a^2 - b}$ are real provided $a^2 \geq b$, which imposes $b \leq 0$ or $b > 0$ and either $a \geq \sqrt{b}$ or $a \leq -\sqrt{b}$. The corresponding eigenfunctions on $x < 0$ are $\phi_{1\pm}(x) = x + a \pm \sqrt{a^2 - b}$. Hence $v_0 = a \pm \sqrt{a^2 - b}$ and $v_1 = 1$. On the whole line, we can get an odd wavefunction if $a = \mp\sqrt{a^2 - b}$, which imposes $b = 0$ and $a < 0$ (resp. $a > 0$) for ϕ_{1+} (resp. ϕ_{1-}). We conclude that the potential

$$V(x) = x^4 - 4a|x|^3 + 4a^2x^2 - 4|x| \tag{2.13}$$

has an energy eigenvalue $E_1^{(-)}$ and associated wavefunction $\psi_1^{(-)}(x)$, given by

$$E_1^{(-)} = -6a, \qquad \psi_1^{(-)}(x) = e^{-\frac{1}{3}|x|^3 + ax^2}x, \tag{2.14}$$

corresponding to a first-excited state.

We may also ask whether an even wavefunction can be obtained for $n = 1$. From (2.5), we directly get that $b \neq 0$ and $a \pm \sqrt{a^2 - b} = -1/b$, where the upper (resp. lower) sign corresponds to ϕ_{1+} (resp. ϕ_{1-}). This implies that a can be expressed in terms of b as $a = -(b^3 + 1)/(2b)$. As a result, $\sqrt{a^2 - b} = (b^3 - 1)/(2b)$ and $a + \sqrt{a^2 - b} = -1/b$ if $b < 0$ or $b > 1$, whereas $\sqrt{a^2 - b} = (1 - b^3)/(2b)$ and $a - \sqrt{a^2 - b} = -1/b$ if $0 < b \leq 1$. In both cases, $E_{1+} = E_{1-} = (2b^3 + 1)/b$. Hence, the potential

$$V(x) = x^4 + \frac{2(b^3 + 1)}{b}|x|^3 + \frac{b^6 + 4b^3 + 1}{b^2}x^2 + 2(b^3 - 1)|x|, \qquad b \neq 0, \tag{2.15}$$

has an energy eigenvalue $E_1^{(+)}$ and associated wavefunction $\psi_1^{(+)}(x)$, given by

$$E_1^{(+)} = \frac{2b^3 + 1}{b}, \qquad \psi_1^{(+)}(x) = e^{-\frac{1}{3}|x|^3 - \frac{b^3+1}{2b}x^2 - b|x|}\left(|x| + \frac{1}{b}\right), \tag{2.16}$$

corresponding to a ground state for $b > 0$ and to a second-excited state for $b < 0$.



Figure 1: The potential $V(x)$, defined in Eq. (3.8), in terms of x for $a = 0$ (black line), $a = +1$ (red line), and $a = -1$ (blue line). In all cases, $b = -1$ and $E_0^{(+)} = 2$.

3 QES Symmetrized Sextic Polynomial Oscillator

Let us now consider the symmetrized sextic polynomial oscillator

$$V(x) = x^6 - u|x|^5 + tx^4 - s|x|^3 + rx^2 - q|x|, \tag{3.1}$$

where q, r, s, t, and u are five real constants, and let us proceed as in Sec. 2. QES solutions of (2.2) with potential (3.1) may be obtained by assuming that

$$\psi_n^{(\epsilon)}(x) = e^{W(x)}\phi_n^{(\epsilon)}(x), \qquad W(x) = -\frac{1}{4}x^4 - a|x|^3 + bx^2 - c|x|, \tag{3.2}$$

where a, b, c are three real constants and $\phi_n^{(\epsilon)}(x)$ is still given by Eq. (2.4), but with Eq. (2.5) replaced by the constraint

$$\begin{aligned} &v_1 + cv_0 = 0 \qquad \text{if } \epsilon = +1, \\ &v_0 = 0 \qquad \text{if } \epsilon = -1. \end{aligned} \tag{3.3}$$

By setting now

$$u = -6a, \qquad t = 9a^2 - 4b, \qquad s = 12ab - 2c, \tag{3.4}$$



Figure 2: The QES ground-state wavefunction $\psi_0^{(+)}(x)$, defined in Eq. (3.9), in terms of x for $a = 0$ (black line), $a = +1$ (red line), and $a = -1$ (blue line). In all cases, $b = -1$ and $E_0^{(+)} = 2$.

we get on $x < 0$ the differential equation

$$\left(-\frac{d^2}{dx^2} + 2(x^3 - 3ax^2 - 2bx - c)\frac{d}{dx} + (r - 4b^2 - 6ac + 3)x^2 + (q - 4bc - 6a)x - c^2 - 2b\right)\phi_n(x) = E_n\phi_n(x), \tag{3.5}$$

which is (a special case of) a generalized Heun equation. Its nth-degree polynomial solutions $\phi_n(x) = \prod_{i=1}^n (x - x_i)$ with distinct roots x_1, x_2, ..., x_n, can be found by applying the functional Bethe ansatz method. Direct application of Theorem 1.1 of Ref. [8] leads to the following conditions

$$\begin{aligned} r &= 4b^2 + 6ac - 2n - 3, \qquad q = -2\sum_{i=1}^n x_i + 6a(n+1) + 4bc, \\ E &= 2\sum_{i=1}^n x_i^2 - 6a\sum_{i=1}^n x_i - 2b(2n+1) - c^2, \end{aligned} \tag{3.6}$$

where the roots x_1, x_2, ..., x_n satisfy the Bethe ansatz equations

$$\sum_{\substack{j=1 \\ j\neq i}}^n \frac{1}{x_i - x_j} - x_i^3 + 3ax_i^2 + 2bx_i + c = 0, \qquad i = 1, 2, \ldots, n. \tag{3.7}$$



Figure 3: The potential $V(x)$, defined in Eq. (3.12), in terms of x for $a = +1$, $b = -1$, $c = -1.949788$ (red line), and $a = -1$, $b = -1$, $c = -0.458984$ (blue line). The corresponding energy is $E_1^{(+)} = 5.80187$ and $E_1^{(+)} = 2.21067$, respectively.

On the resulting $\phi_n(x)$, it then remains to impose constraint (3.3).

The $n = 0$ is straightforwardly solved. As for the quartic oscillator, there is no odd eigenfunction, but there is an even one provided $c = 0$. Hence, the potential

$$V(x) = x^6 + 6a|x|^5 + (9a^2 - 4b)x^4 - 12ab|x|^3 + (4b^2 - 3)x^2 - 6a|x| \tag{3.8}$$

has an energy eigenvalue $E_0^{(+)}$ and associated wavefunction $\psi_0^{(+)}(x)$, given by

$$E_0^{(+)} = -2b, \qquad \psi_0^{(+)}(x) = e^{-\frac{1}{4}x^4 - a|x|^3 + bx^2}, \tag{3.9}$$

corresponding to a ground state.

For $n = 1$, there is a single Bethe ansatz equation to be solved, namely $x_1^3 - 3ax_1^2 - 2bx_1 - c = 0$. Since $v_0 = -x_1$ and $v_1 = 1$, we can get an even or odd wavefucntion by appropriately choosing the parameters. The function is odd if $x_1 = 0$, implying $c = 0$. Hence the potential

$$V(x) = x^6 + 6a|x|^5 + (9a^2 - 4b)x^4 - 12ab|x|^3 + (4b^2 - 5)x^2 - 12a|x| \tag{3.10}$$

has an energy eigenvalue $E_1^{(-)}$ and associated wavefunction $\psi_i^{(-)}(x)$, given by

$$E_1^{(-)} = -6b, \qquad \psi_1^{(-)}(x) = e^{-\frac{1}{4}x^4 - a|x|^3 + bx^2} x, \tag{3.11}$$



Figure 4: The QES second-excited state wavefunction $\psi_1^{(+)}(x)$, defined in Eq. (3.13), in terms of x for $a = +1$, $b = -1$, $c = -1.949788$ (red line), and $a = -1$, $b = -1$, $c = -0.458984$ (blue line). The corresponding energy is $E_1^{(+)} = 5.80187$ and $E_1^{(+)} = 2.21067$, respectively.

corresponding to a first-excited state. On the other hand, the function is even if $x_1 = 1/c$, where c must be a nonvanishing solution of the quartic equation $c^4 + 2bc^2 + 3ac - 1 = 0$. Then, for such a c value, the potential

$$V(x) = x^6 + 6a|x|^5 + (9a^2 - 4b)x^4 - (12ab - 2c)|x|^3 + (4b^2 + 6ac - 5)x^2 - \left(12a + 4bc - \frac{2}{c}\right)|x| \tag{3.12}$$

admits $E_1^{(+)}$ and $\psi_1^{(+)}(x)$, given by

$$E_1^{(+)} = -6b - c^2 - \frac{6a}{c} + \frac{2}{c^2}, \qquad \psi_1^{(+)}(x) = e^{-\frac{1}{4}x^4 - a|x|^3 + bx^2 - c|x|}\left(|x| + \frac{1}{c}\right), \tag{3.13}$$

representing a ground state for $c > 0$ and a second-excited state for $c < 0$.

For $n = 2$ and $\epsilon = -1$, we get $x_1 = 0$ and $x_2 = 1/c$, where c is a nonvanishing

solution of $2c^4+2bc^2+3ac-1=0$. This yields

$$\begin{aligned} V(x) &= x^6+6a|x|^5+(9a^2-4b)x^4-(12ab-2c)|x|^3+(4b^2+6ac-7)x^2 \\ &\quad -\left(18a+4bc-\frac{2}{c}\right)|x|, \\ E_2^{(-)} &= -10b-c^2-\frac{6a}{c}+\frac{2}{c^2}, \\ \psi_2^{(-)}(x) &= e^{-\frac{1}{4}x^4-a|x|^3+bx^2-c|x|}x\left(|x|+\frac{1}{c}\right), \end{aligned} \tag{3.14}$$

where $\psi_2^{(-)}(x)$ corresponds to a first-excited (resp. third-excited) state for $c>0$ (resp. $c<0$).

Furthermore, for $n=2$ and $\epsilon=+1$, we obtain

$$\begin{aligned} V(x) &= x^6+6a|x|^5+(9a^2-4b)x^4-(12ab-2c)|x|^3+(4b^2+6ac-7)x^2 \\ &\quad -(18a+4bc-2x_1-2x_2)|x|, \\ E_2^{(+)} &= -10b-c^2-6a(x_1+x_2)+2(x_1^2+x_2^2), \\ \psi_2^{(-)}(x) &= e^{-\frac{1}{4}x^4-a|x|^3+bx^2-c|x|}(|x|+x_1)(|x|+x_2), \end{aligned} \tag{3.15}$$

where x_1 and x_2 are solutions different from zero and $1/c$ of the set of equations

$$x_1^3-3ax_1^2-2bx_1-c=-(x_2^3-3ax_2^2-2bx_2-c)=\frac{1}{x_1-x_2}, \qquad x_1+x_2=cx_1x_2. \tag{3.16}$$

The set of QES symmetrized sextic polynomial oscillators that we have constructed here can be separated into two categories.

(i) The symmetrized sextic oscillators whose known wavefunction is of 'natural' parity, i.e., such that $\epsilon=(-1)^n$. These include those associated with the cases $(n,\epsilon)=(0,+1)$, $(1,-1)$, and $(2,+1)$, considered above. It is straightforward to see that they comprise as special cases the well-known (analytic) sextic oscillators of Ref. [10]. This is obvious if we set $a=0$ in Eqs. (3.8)–(3.11), corresponding to $(n,\epsilon)=(0,+1)$ and $(1,-1)$. In the case of Eq. (3.15) for $(n,\epsilon)=(2,+1)$, on assuming $a=c=x_1+x_2=0$, the set of equations (3.16) reduces to a single equation $2x_1^4-4bx_1^2-1=0$, leading to $(x_1^2)_\pm=\frac{1}{2}(2b\pm\sqrt{4b^2+2})$. As a consequence, $E_{2\pm}^{(+)}=-6b\pm2\sqrt{4b^2+2}$ and $\psi_{2\pm}^{(+)}(x)=\exp(-\frac{1}{4}x^4+bx^2)(x^2-b\mp\frac{1}{2}\sqrt{4b^2+2})$, in agreement with [10].

(ii) The symmetrized sextic oscillators whose known wavefunction is of 'unnatural' parity, i.e., such that $\epsilon = (-1)^{n+1}$. These include those associated with the cases $(n, \epsilon) = (1, +1)$ and $(2, -1)$, considered above. Such symmetrized oscillators do not comprise any analytic sextic oscillator because the $c \neq 0$ condition always ensures the nonvanishing of an $|x|^3$ term in $V(x)$.

In Figs. 1 and 2, some examples of potential and associated wavefunction are displayed for the case $(n, \epsilon) = (0, +1)$, belonging to the first category. They are compared with the corresponding results for the analytic potential with $a = 0$. In Figs. 3 and 4, some examples of potential and associated wavefunction are plotted for the case $(n, \epsilon) = (1, +1)$, belonging to the second category.

4 Conclusion

In the present work, we have shown that the previously introduced QES symmetrized quartic polynomial oscillator admits an sl(2,$\mathbb{R}$) algebraization and we have used the latter to provide some simple examples of bound-state wavefunctions at certain couplings and energies.

We have then proposed a new symmetrized sextic polynomial oscillator and we have proved its QES nature by deriving some exact, closed-form solutions of the corresponding Schrödinger equation by resorting to the functional Bethe ansatz method. Furthermore, we have shown that such symmetrized sextic oscillators can be separated into two categories: a first set generalizing the well-known analytic sextic oscillators, which form a subset, and a second set of novel QES potentials with no counterpart in the class of analytic sextic oscillators.

Trying to extend other known QES analytic potentials by allowing some additional analyticity-violating terms would provide an interesting topic for future investigation.

References

[1] H. Ciftci, R. I. Hall, N. Saad and E. Dogu, *J. Phys. A* **43**, 415206 (2010), arXiv:1009.0464.

[2] A. V. Turbiner, *Commun. Math. Phys.* **118**, 467 (1988).

[3] A. G. Ushveridze, *Quasi-Exactly Solvable Models in Quantum Mechanics* (IOP, Bristol, 1994).

[4] A. González-López, N. Kamran and P. J. Olver, *Commun. Math. Phys.* **153**, 117 (1993).

[5] A. V. Turbiner, *Phys. Reports* (in press), arXiv:1603.02992.

[6] M. Gaudin, *La Fonction d'Onde de Bethe* (Masson, Paris, 1983).

[7] C.-L. Ho, *Ann. Phys. (N. Y.)* **323**, 2241 (2008), arXiv:0801.0944.

[8] Y.-Z. Zhang, *J. Phys. A* **45**, 065206 (2012), arXiv:1107.5090.

[9] P. G. L. Leach, *J. Math. Phys.* **25**, 2974 (1984).

[10] A. V. Turbiner and A. G. Ushveridze, *Phys. Lett. A* **126**, 181 (1987).

[11] M. Znojil, *Phys. Lett. A* **381**, 1414 (2016), arXiv:1602.07088.

[12] M. Znojil, *Mod. Phys. Lett. A* **31**, 1650088 (2016), arXiv:1603.09483.

[13] R. Sasaki and M. Znojil, arXiv:1605.07310.

[14] N. Mohammedi and T. R. Morris, arXiv:1606.06637.

[15] A. Ronveaux, *Heun Differential Equations* (Oxford University Press, Oxford, 1995).